Health Link

Healthy living after treatment for childhood cancer

Colorectal Cancer Version 3.0 - 10/08

Colorectal Cancer Following Treatment for Childhood Cancer: Are You at Risk?

You have successfully been treated for cancer during childhood or adolescence and are now moving forward with your life - so the last thing you want to be reminded about is the risk of developing another cancer during adulthood. For a variety of reasons, the risk of cancer increases for everyone as they age. Depending on the specific treatment you received for childhood cancer, you may be at increased risk for developing colorectal cancer (cancer of the colon or rectum). It is important to understand that risk, so that you can take steps to protect your health.

What is colorectal cancer?

Colorectal cancer is a type of cancer that occurs in the colon (large intestine) or the rectum (the last several inches of the large intestine). Colorectal cancer is the second leading cause of cancer deaths in the United States. Many of these deaths happen because the cancers are found too late to be cured. If colorectal cancer is found early enough, it can usually be cured.

What are the risk factors?

Several studies have shown that people who were treated with radiation to the abdomen, pelvis, or spine in moderate to high doses during childhood, adolescence, or young adulthood have an increased risk of developing colorectal cancer. It is therefore important for you to obtain your radiation treatment records so that you know how much radiation you received.

Other known risk factors for developing colorectal cancer include:

- Having had colorectal cancer or large intestinal polyps in the past
- Having a close relative (brother, sister, parent or child) who has had colorectal cancer before age 60
- Having ulcerative colitis or Crohn's disease
- Having a hereditary colon cancer syndrome (such as familial adenomatous polyposis)

What are the signs of colorectal cancer?

Most colorectal cancers begin as a polyp. A polyp starts as a small, harmless growth in the wall of the colon or rectum. However, as a polyp gets larger, it can develop into a cancer that grows and spreads. During the early stage of colorectal cancer, there are



Health Link

Healthy living after treatment for childhood cancer

Colorectal Cancer Version 3.0 - 10/08

rarely any outward signs or symptoms to alert you or your healthcare provider that cancer is present. This is why screening is so important. Once the cancer has become more advanced, the following signs may be evident. If you have any of these signs, you should see your healthcare provider immediately:

- Bleeding from your rectum
- Blood in your stool or in the toilet after you have a bowel movement
- A change in the shape of your stool
- Cramping pain in your lower stomach
- A feeling of discomfort or an urge to have a bowel movement when there is no need to have one
- A change in the normal frequency of your bowel movements

Other conditions can cause these same symptoms. You should be evaluated by your healthcare provider to find out the reason for your symptoms.

When is colorectal cancer likely to occur?

In the general population, colorectal cancer is most likely to occur between the ages of 50 and 65. In cancer survivors who were treated with abdominal, pelvic, or spinal radiation, it may occur earlier. The risk begins to increase around 10 years after the radiation.

What can I do to protect my health?

Most people who received radiation therapy to the abdomen, pelvis, or spine will **NOT** develop colorectal cancer. However, if you received this type of radiation, it is important to understand that the risk **IS** higher for you than it is for other people your age who never received radiation. So, the best way for you to protect your health is by taking steps to closely monitor your colon. That way, if a cancer develops, it can be detected in its earliest stages, when treatment is most effective.

What monitoring is recommended?

If you were treated with radiation therapy to the abdomen, pelvis, or spine at doses of 30 Gy (3000 cGy/rads) or higher* during childhood, adolescence, or young adulthood, you should have a colonoscopy at least every five years, starting at the age of 35, or 10 years after radiation (whichever occurs last).

*Note: If you received total body irradiation (TBI), your TBI dose should be added to this total.



Colorectal Cancer Version 3.0 - 10/08

What is a colonoscopy?

Before you have this test, you will be given a medicine to make you feel relaxed and sleepy. A thin, flexible tube connected to a video camera is then inserted into your rectum and slowly guided into your colon. The doctor is able to examine your entire colon, and any polyps or growths can be removed through the tube during the exam. Colonoscopy may be uncomfortable, but it is usually not painful.

What if I had a low dose of radiation to these areas (or had total body irradiation)?

For those treated with lower doses of radiation therapy to the abdomen, pelvis, or spine (less than 30 Gy or 3000 cGy/rads), or for those who had total body radiation, there still may be an increased risk, but there are no scientific studies currently available to confirm this. You should have a discussion with your healthcare provider and make a decision about whether or not monitoring with colonoscopy should be started earlier for you than for people who never had radiation.

Is there anything else that I can do to minimize the risk?

Yes, most definitely! Studies have shown that the following lifestyle changes reduce the risk of colorectal cancer and will help you stay as healthy as possible:

Eat a variety of healthful foods, with an emphasis on grains, fruits and vegetables

- Eat five or more servings of a variety of vegetables and fruits each day
- Choose whole grains in preference to processed (refined) grains and sugars
- Limit consumption of red meats, especially processed meats (such as hot dogs or bologna) and those high in fat
- Choose foods that maintain a healthy weight

Adopt a physically active lifestyle.

- Engage in at least moderate physical activity (such as brisk walking) for 30 minutes or more on five or more days of the week
- Engaging in 45 minutes or more of moderate to vigorous activity (activities such as running, in which you are not able to carry on a conversation without needing to catch your breath) on five or more days per week may further reduce your risk of colon cancer



Health Link

Healthy living after treatment for childhood cancer

Colorectal Cancer Version 3.0 - 10/08

Written by Kevin Oeffinger MD, Division of Pediatrics, Memorial Sloan-Kettering Cancer Center, New York, New York.

Reviewed by Jacqueline Casillas MD, Wendy Landier CPNP, and Joan Darling PhD.

Additional health information for childhood cancer survivors is available at www.survivorshipquidelines.org

Note: Throughout this *Health Links* series, the term "childhood cancer" is used to designate pediatric cancers that may occur during childhood, adolescence, or young adulthood. *Health Links* are designed to provide health information for survivors of pediatric cancer, regardless of whether the cancer occurred during childhood, adolescence, or young adulthood.

Disclaimer and Notice of Proprietary Rights

Introduction to Late Effects Guidelines and Health Links: The "Long-Term Follow-Up Guidelines for Survivors of Childhood, Adolescent, and Young Adult Cancers" and accompanying "Health Links" were developed by the Children's Oncology Group as a collaborative effort of the Late Effects Committee and Nursing Discipline and are maintained and updated by the Children's Oncology Group's Long-Term Follow-Up Guidelines Core Committee and its associated Task Forces. For Informational Purposes Only: The information and contents of each document or series of documents made available from by the Children's Oncology Group relating to late effects of cancer treatment and care or containing the title "Long-Term Follow-Up Guidelines for Survivors of Childhood, Adolescent, and Young Adult Cancers" or the title "Health Link", whether available in print or electronic format (including any digital format, e-mail transmission, or download from the website), shall be known hereinafter as "Informational Content". All Informational Content is for informational purposes only. The Informational Content is not intended to substitute for medical advice, medical care, diagnosis or treatment obtained from a physician or health care provider.

To cancer patients (if children, their parents or legal guardians): Please seek the advice of a physician or other qualified health provider with any questions you may have regarding a medical condition and do not rely on the Informational Content. The Children's Oncology Group is a research organization and does not provide individualized medical care or treatment.

To physicians and other healthcare providers: The Informational Content is not intended to replace your independent clinical judgment, medical advice, or to exclude other legitimate criteria for screening, health counseling, or intervention for specific complications of childhood cancer treatment. Neither is the Informational Content intended to exclude other reasonable alternative follow-up procedures. The Informational Content is provided as a courtesy, but not intended as a sole source of guidance in the evaluation of childhood cancer survivors. The Children's Oncology Group recognizes that specific patient care decisions are the prerogative of the patient, family, and healthcare provider.

No endorsement of any specific tests, products, or procedures is made by Informational Content, the Children's Oncology Group, or affiliated party or member of the Children's Oncology Group.

No Claim to Accuracy or Completeness: While the Children's Oncology Group has made every attempt to assure that the Informational Content is accurate and complete as of the date of publication, no warranty or representation, express or implied, is made as to the accuracy, reliability, completeness, relevance, or timeliness of such Informational Content.

No Liability on Part of Children's Oncology Group and Related Parties/ Agreement to Indenmify and Hold Harmless the Children's Oncology Group and Related Parties:

No liability is assumed by the Children's Oncology Group or any affiliated party or member thereof for damage resulting from the use, review, or access of the Informational Content. You agree to the following terms of indemnification: (i) "Indemnified Parties" include authors and contributors to the Informational Content, all officers, directors, representatives, employees, agents, and members of the Children's Oncology Group and affiliated organizations; (ii) by using, reviewing, or accessing the Informational Content, you agree, at your own expense, to indemnify, defend and hold harmless Indemnified Parties from any and all losses, liabilities, or damages (including attorneys' fees and costs) resulting from any and all claims, causes of action, suits, proceedings, or demands related to or arising out of use, review or access of the Informational Content.

Proprietary Rights: The Informational Content is subject to protection under the copyright law and other intellectual property law in the United States and worldwide. The Children's Oncology Group retains excursive copyright and other right, title, and interest to the Informational Content and claims all intellectual property rights available under law. You hereby agree to help the Children's Oncology Group secure all copyright and intellectual property rights for the benefit of the Children's Oncology Group by taking additional action at a later time, action which could include signing consents and legal documents and limiting dissemination or reproduction of Informational Content.

